

Roll No. ....

**DD-763**

**M. A./M. Sc. (Fourth Semester)**

**EXAMINATION, 2020**

**MATHEMATICS**

**Paper Second**

**(Partial Differential Equations and Mechanics)**

*Time : Three Hours*

*Maximum Marks : 80*

**Note :** All questions are compulsory. Attempt any *two* parts from each question. All questions carry equal marks.

**Unit—I**

1. (a) State and prove local existence theorem for first order non-linear PDE.
- (b) Prove that the function  $x(.)$  solves the system of Euler-Lagrange's equations :

$$-\frac{d}{ds}(D_q L(\dot{x}(s), x(s))) + D_x L(\dot{x}(s), x(s)) = 0$$

$(0 \leq s \leq t)$

- (c) State and prove Lax-Oleinik's formula.

**P. T. O.**

## Unit—II

2. (a) Using the method of separation of variables solve the heat equation :

$$u_t - \Delta u = 0 \text{ in } U \times (0, \infty)$$

$$u = 0 \text{ on } \partial U \times [0, \infty)$$

$$u = g \text{ on } U \times \{t = 0\}$$

where  $g : U \rightarrow \mathbb{R}$  is given.

- (b) Derive Barenblott's solution to the porous medium equation using similarity under scaling.  
(c) Explain Hodograph transform.

## Unit—III

3. (a) Suppose  $k, l : \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions, that  $l$  grows at most linearly and that  $k$  grows at least quadratically. Assume also there exists a unique point  $y_0 \in \mathbb{R}$  such that :

$$k(y_0) = \min_{y \in \mathbb{R}} k(y)$$

Then prove that :

$$\lim_{\epsilon \rightarrow 0} \frac{\int_{-\infty}^{\infty} l(y) e^{-\frac{k(y)}{\epsilon}} dy}{\int_{-\infty}^{\infty} e^{-\frac{k(y)}{\epsilon}} dy} = l(y_0).$$

- (b) Discuss unperturbed PDE :

$$\operatorname{div}(u b) = \delta_0 \text{ in } \mathbb{R}^2$$

where  $\delta_0$  is the Dirac measure  $\mathbb{R}^2$  giving unit mass to the point  $O$ .

- (c) Solve the wave equation using stationary phase method.

## Unit—IV

4. (a) Derive mathematical expressions for Hamilton's principle.  
(b) Derive Whittaker's equations.  
(c) State and prove Lee Hwa-Chung's theorem.

## Unit—V

5. (a) Derive Hamilton-Jacobi's equations.  
(b) Prove that the Lagrange bracket is invariant under canonical transformation.  
(c) Using Poisson brackets relation, show that the following transformation is canonical if  $ad - bc = 1$  :

$$Q = aq + bp$$

$$P = cq + dp$$